The new multijet trigger for $H + W/Z$

Petar Maksimović

for

J. Huth, W. Riegler, T. Rodrigo, J. Troconiz, J. Valls, R. Vilar

Trigger and Datasets Working Group, Jan 14, 1999

In this talk:

1. *Old* multijet trigger – and what's wrong with it
2. New multijet trigger basics
3. Run2Trs in action
4. What to do at Level 1?

Next time:

1. Use of tracking information: XFT and SVT
2. b-tagging at Level 3? (SECVTX, JPB, 1 or 2 tags...)

Old multijet trigger – and what's wrong with it

Run I multijet trigger:

- Designed for $t\bar{t} \to 6$ jets
- 4 jets, $E_T > 15$ GeV (10 GeV in Run Ia)
- $\sum E_T > 125$ GeV ($\sum E_T > 100$ in Run Ia)

Problems for Higgs: low efficiency

- $\sum E_T > 125$ is too high
- $N \geq 4$ jets too restrictive (in trigger)

Improving efficiency for $H + W/Z$



New multijet trigger – basics

1. relax number of jets to 3
2. relax $\sum E_T$ cut down to 70 GeV

New multijet trigger – basics

Must cut the rate elsewhere.

New cuts:

1. **at least one displaced track(s) at Level 2**
2. **two b-tags (SECVTX) tags at Level 3**
   (can run offline code at Level 3)

Figures of merit

- **signal efficiency**
  use $H + W/Z$ Pythia Monte Carlo
  ($m_H = 110$ GeV)
- **Level 1, Level 2, Level 3 rates**
  use Jet 20 data
  (since $E_T > 20$ GeV for leading jet)
- $\sigma(\mathrm{Jet20}) = 7.5\mu b$

L = 10$^{32}cm^{-2}s^{-1}$

Run2Trs in action

- Previously, we used Run I TRGSIM/TRVIEW, for Higgs MC and used real trigger info from Jet 20
- Now we apply Run2Trs to both

  $\Rightarrow$ some Jet 20 variables are smeared

  (e.g. $E_T(1) > 20$ GeV!)
- Also, now we have Level 1 $\sum E_T$