

First glance at $B^0 \rightarrow \eta' K^0$ time-dependent CPV analysis

Stefano Lacaprara

INFN Padova

23rd B2GM, Physics session
KEK, February 1st 2016

- 1 Introduction and motivations
- 2 $B^0 \rightarrow \eta'(\rightarrow \eta_{\gamma\gamma} \pi^+ \pi^-)(K_S^0 \rightarrow \pi^+ \pi^-)$
 - Background
 - $B^0 \rightarrow \eta'(\rightarrow \eta_{\gamma\gamma} \pi^+ \pi^-)(K_S^0 \rightarrow \pi^0 \pi^0)$
- 3 $B^0 \rightarrow \eta'(\rightarrow \eta_{3\pi} \pi^+ \pi^-)(K_S^0 \rightarrow \pi^+ \pi^-)$
 - Background
 - $B^0 \rightarrow \eta'(\rightarrow \eta_{3\pi} \pi^+ \pi^-)(K_S^0 \rightarrow \pi^0 \pi^0)$
- 4 Summary and outlook

1 Introduction and motivations



- 2 $B^0 \rightarrow \eta'(\rightarrow \eta_{\gamma\gamma} \pi^+ \pi^-)(K_S^0 \rightarrow \pi^+ \pi^-)$
- Background
 - $B^0 \rightarrow \eta'(\rightarrow \eta_{\gamma\gamma} \pi^+ \pi^-)(K_S^0 \rightarrow \pi^0 \pi^0)$

- 3 $B^0 \rightarrow \eta'(\rightarrow \eta_{3\pi} \pi^+ \pi^-)(K_S^0 \rightarrow \pi^+ \pi^-)$
- Background
 - $B^0 \rightarrow \eta'(\rightarrow \eta_{3\pi} \pi^+ \pi^-)(K_S^0 \rightarrow \pi^0 \pi^0)$

4 Summary and outlook

A sensitivity study for Time-Dependent CP violation analysis in the $B^0 \rightarrow \eta' K^0$ channel, a charmless $b \rightarrow sq\bar{q}$ decay

- CP asymmetry from time-dependent decay rate into CP eigenstates;
- $S_{\eta' K^0} = \sin 2\phi_1^{eff}$ tightly related to $\sin 2\phi_1$
 - ▶ identical if only penguin diagram were present;
 - ▶ QCD factorization $\Delta S_{\eta' K^0} \in [-0.03, 0.03]$
 - ▶ new physics can enter in the loop, shifting $\Delta S_{\eta' K^0}$ more than SM expectation
- Similar to $B^0 \rightarrow \phi K^0$, see work by A. Gaz
 - ▶ more complex final state;
 - ▶ η' is a pseudo-scalar (ϕ a vector);
 - ▶ large BR: $\sim 6.6 \cdot 10^{-5}$ ($\sim 10 \times \text{BR}(B^0 \rightarrow \phi K^0)$)^[CLEO(1998)]
 - ★ constructive interference between penguin diagrams
 - ▶ actual uncertainties $\sigma_{stat} = 0.07, \sigma_{syst} = 0.03$ ^[Belle(2014)]
 - ▶ projected for 50 ab^{-1} $\sigma_{stat} = 0.008, \sigma_{syst} = 0.008$ ^[Urquijo(2015)]
 - ▶ no competition from LHCb (neutrals);



- Channel have been analyzed in B-factory ^[BABAR(2009), Belle(2007), Belle(2014)];
- uncertainties are mostly statistical (~ 3500 events for all final states);
- quasi-two body approach;
- many decay channels available $B^0 \rightarrow \eta' K_S^0$
 - $\eta' \rightarrow \rho(\rightarrow \pi^+ \pi^-) \gamma$; BR: 29% not yet: Hulya Atmacan (METU) is interested in working on this
 - $\eta' \rightarrow \eta \pi^+ \pi^-$; BR: 43%
 - $\eta_{\gamma\gamma}: \eta \rightarrow \gamma\gamma$; BR: 40%
 - $\eta_{3\pi}: \eta \rightarrow \pi^+ \pi^- \pi^0$; BR: 23%
 - $K_S^0 \rightarrow \pi^+ \pi^-$, $K_S^0 \rightarrow \pi^0 \pi^0$,
 - Total BR($B_0 \rightarrow \eta'(\rightarrow_{\eta_{\gamma\gamma}/\eta_{3\pi}} \pi^+ \pi^-) K_S^0$) = 27% today
- $B^0 \rightarrow \eta' K_L^0$ not yet studied
- Complex final state, neutrals, large combinatorics;
- Release used rel-00-05-03
- Signal: MC5 BGx0 (some private production); Background MC5 BGx1

- 1 Introduction and motivations
- 2 $B^0 \rightarrow \eta'(\rightarrow \eta_{\gamma\gamma} \pi^+ \pi^-)(K_S^0 \rightarrow \pi^+ \pi^-)$
 - Background
 - $B^0 \rightarrow \eta'(\rightarrow \eta_{\gamma\gamma} \pi^+ \pi^-)(K_S^0 \rightarrow \pi^0 \pi^0)$
- 3 $B^0 \rightarrow \eta'(\rightarrow \eta_{3\pi} \pi^+ \pi^-)(K_S^0 \rightarrow \pi^+ \pi^-)$
 - Background
 - $B^0 \rightarrow \eta'(\rightarrow \eta_{3\pi} \pi^+ \pi^-)(K_S^0 \rightarrow \pi^0 \pi^0)$
- 4 Summary and outlook

$$B^0 \rightarrow \eta'(\rightarrow \eta_{\gamma\gamma} \pi^+ \pi^-)(K_S^0 \rightarrow \pi^+ \pi^-)$$

- **skim** at least one decay chain have been reconstructed
 - ▶ w/o vertex or mass fit and with loose selections (as for the background)
 - ▶ Using the same skim for all decay channels: run only once on background
- **pre-selection** $N \geq 1$ decay chain, with vertex/mass constraint
- **selection** cuts on reconstructed quantities


good candidate selection:

- | | |
|--|---|
| ▶ $M_{bc} > 5.25$ GeV; | ▶ $\text{PID}_{\pi}(\pi^{\pm}) > 0.2$; Should use $\mathcal{L}$ -ratio instead |
| ▶ $ \Delta E < 0.1$ GeV; | ▶ $d_0(\pi^{\pm}) < 0.08$ mm; |
| ▶ $M(\eta_{\gamma\gamma}) \in [0.45, 0.57]$ GeV; | ▶ $z_0(\pi^{\pm}) < 0.1$ mm; |
| ▶ $M(\eta') \in [0.93, 0.98]$ GeV; | ▶ $N \text{ hits}_{\text{PXD}}(\pi^{\pm}) > 1$ |
| ▶ $M(K_S^0 \rightarrow \pi^+ \pi^-) \in [0.48, 0.52]$ GeV; | ▶ $\text{P-value}_{\text{vtx}}(B_0, \eta', K_S^0) > 1 \cdot 10^{-5}$ |


Best candidate

if $N_{\text{cands}} > 1$, select candidate with highest $\text{P-value}_{\text{vtx}}(B_0, \eta', \eta, K_S^0)$

Distributions: all/good candidates




- combinatorics small
- some true cand lost (mostly due to $PID(\pi)$ and $N \text{ hits}_{PXD}(\pi^\pm) > 1$);
- almost all good candidate(s) are true;
- $M_{\eta'}$ has a lower tail;





All best candidates are true: SXF is negligible

Efficiency and Δt resolution

Cands multiplicity



Efficiency %	
skim	57.1
preselection	48.1
good cands	33.1
MC true	31.9
SXF	1.2









Flavour tagging is random,
with $\epsilon(1 - 2w)^2 = 32\%$ dilution

$$B^0 \rightarrow \eta'(\rightarrow \eta_{\gamma\gamma} \pi^+ \pi^-)(K_S^0 \rightarrow \pi^+ \pi^-)$$

- Background MC sample **BGx1**, single skim for all channels;
- Using only a fraction: **L=30 fb⁻¹**;
 - ▶ need to process the full MC5 dataset to increase # of selected events;
 - ▶ Need to reduce the skim output!
- **NB No cut (yet) on continuum discriminating variable**
 - ▶ still learning how to use it
 - ▶ Alessandro showed some issue in $B^0 \rightarrow \phi K^0$ presentation on 2015/12/11

Sample	# Ev (M)	Skim (M)	ϵ_{skim}	pre-sel	sel	ϵ_{sel}
$u\bar{u}$	48.15	1.023	$2.13 \cdot 10^{-2}$	8196	93	$1.96 \cdot 10^{-6}$
$d\bar{d}$	13.03	0.274	$2.27 \cdot 10^{-2}$	2333	37	$2.84 \cdot 10^{-6}$
$s\bar{s}$	11.49	0.238	$2.07 \cdot 10^{-2}$	2334	18	$1.57 \cdot 10^{-6}$
$c\bar{c}$	38.87	1.127	$2.90 \cdot 10^{-2}$	9911	123	$1.09 \cdot 10^{-6}$
total	111.54	2.662	$2.39 \cdot 10^{-2}$	22774	271	$2.43 \cdot 10^{-6}$


$$B^0 \rightarrow \eta'(\eta(\gamma\gamma) \pi^+\pi^-) K_S^0(\pi^+\pi^-)$$



$$B^0 \rightarrow \eta'(\rightarrow \eta_{\gamma\gamma} \pi^+ \pi^-)(K_S^0 \rightarrow \pi^0 \pi^0)$$

- Similar selection as in $K_S^0 \rightarrow \pi^+ \pi^-$ case, taking into account the more difficult π^0 reconstruction

- $\Delta E \in [-0.15, 0.25]$ GeV; (< 0.1 for $K_S^0 \rightarrow \pi^+ \pi^-$)
- $M(\pi^0) \in [0.1, 0.15]$ GeV;
- $M(K_S^0 \rightarrow \pi^0 \pi^0) \in [0.42, 0.52]$ GeV;
- (full list in backup)



- combinatorics larger but still manageable (~ 5 cand's/ev)
- lower tails for $M_{\pi^0, K_S^0, \eta}$;
- SXF still small

Efficiency %	
skim	40.5
preselection	36.1
good cand's	15.6
MC true	13.5
SXF	2.2

$$\epsilon(K_S^{00}) \sim 1/2 \epsilon(K_S^{+-})$$

- 1 Introduction and motivations
- 2 $B^0 \rightarrow \eta'(\rightarrow \eta_{\gamma\gamma} \pi^+ \pi^-)(K_S^0 \rightarrow \pi^+ \pi^-)$
 - Background
 - $B^0 \rightarrow \eta'(\rightarrow \eta_{\gamma\gamma} \pi^+ \pi^-)(K_S^0 \rightarrow \pi^0 \pi^0)$
- 3 $B^0 \rightarrow \eta'(\rightarrow \eta_{3\pi} \pi^+ \pi^-)(K_S^0 \rightarrow \pi^+ \pi^-)$
 - Background
 - $B^0 \rightarrow \eta'(\rightarrow \eta_{3\pi} \pi^+ \pi^-)(K_S^0 \rightarrow \pi^0 \pi^0)$
- 4 Summary and outlook

$$B^0 \rightarrow \eta'(\rightarrow \eta_{3\pi} \pi^+ \pi^-)(K_S^0 \rightarrow \pi^+ \pi^-)$$

Selections $\eta_{3\pi}$







- skim, preselection as for the $\eta_{\gamma\gamma}$ channel

good candidate selection:


- $M_{bc} > 5.25$ GeV;
- $\Delta E \in [-0.15, 0.15]$ GeV;
- $M(\eta_{3\pi}) \in [0.52, 0.57]$ GeV;
- $M(\eta') \in [0.93, 0.98]$ GeV;
- $M(\pi^0) \in [0.1, 0.15]$ GeV;
- $M(K_S^0 \rightarrow \pi^+ \pi^-) \in [0.48, 0.52]$ GeV;
- $\text{PID}_{\pi}(\pi^{\pm}) > 0.2$;
- $d_0(\pi^{\pm}) < 0.08$ mm;
- $z_0(\pi^{\pm}) < 0.15$ mm;
- $N \text{ hits}_{PXD}(\pi^{\pm}) > 1$
- $\text{P-value}_{\text{vtx}}(B_0, \eta', \eta, K_S^0) > 1 \cdot 10^{-5}$

Best candidate

if $N_{\text{cands}} > 1$, select candidate with highest $\text{P-value}_{\text{vtx}}(B_0, \eta', \eta, K_S^0)$




- Combinatorics is huge ($6\pi^\pm$);
- same problem with π^0 ;
- η' and $\eta_{3\pi}$ well reconstructed for good candidates
- non negligible tails for MC true π^0 , η' and $\eta_{3\pi}$
- some 20% of good candidates are false



Sizeable SXF $\sim 20\%$, is smarter best cand selection possible?

Efficiency and Δt resolution



Better resolution ($4\pi^\pm$)

Efficiency %

skim	57.3
preselection	54.8 (?)
good cands	19.1
MC true	15.3
SXF	3.8







- Lower efficiency: only due to π^0 ?
- pre-sel almost 100% on skim? Combinatorics? to be investigated
- Got SWAP problem during event selection on skims, only fraction of skim processed

$$B^0 \rightarrow \eta'(\rightarrow \eta_{3\pi} \pi^+ \pi^-)(K_S^0 \rightarrow \pi^+ \pi^-)$$

- Background MC sample **BGx1**;
- $L=30 \text{ fb}^{-1}$;
- **really** need to process the full MC5 dataset to increment statistics of selected events;
 - ▶ NB No cut (yet) on continuum discriminating variable
- larger contribute from $c\bar{c}$
- reduction is larger ($\sim 10x$) than for $\eta_{\gamma\gamma}$ channel (but ϵ as well)



Sample	# Ev (M)	Skim (M)	ϵ_{skim}	pre-sel	sel	ϵ_{sel}
$u\bar{u}$	48.15	1.023	$2.13 \cdot 10^{-2}$	24037	14	$2.91 \cdot 10^{-7}$
$d\bar{d}$	13.03	0.274	$2.27 \cdot 10^{-2}$	6552	6	$1.25 \cdot 10^{-7}$
$s\bar{s}$	11.49	0.238	$2.07 \cdot 10^{-2}$	9686	5	$1.04 \cdot 10^{-7}$
$c\bar{c}$	38.87	1.127	$2.90 \cdot 10^{-2}$	56653	40	$8.31 \cdot 10^{-7}$
total	111.54	2.662	$2.39 \cdot 10^{-2}$	96928	271	$1.35 \cdot 10^{-7}$

$$B^0 \rightarrow \eta'(\eta(\pi^+\pi^-\pi^0) \pi^+\pi^-) K_S^0(\pi^+\pi^-)$$



$$B^0 \rightarrow \eta'(\rightarrow \eta_{3\pi} \pi^+ \pi^-)(K_S^0 \rightarrow \pi^0 \pi^0)$$

- Channel not used in Belle
- Selection $\sim$ as before
 - $\Delta E \in [-0.15, 0.25]$ GeV; (< 0.15 for $K_S^0 \rightarrow \pi^+ \pi^-$)
 - (full list in backup)



Private production

- combinatorics is even larger (~ 30 candidates per event)
- Sizeable SXF $\sim 30\%$, smarter best cand selection is possible
- $\epsilon(K_S^{00}) \sim 1/2 \epsilon(K_S^{+-})$

Efficiency %

preselection	35.5
good cand's	10.1
best cand	5.96
SXF	3.8

- 1 Introduction and motivations
- 2 $B^0 \rightarrow \eta'(\rightarrow \eta_{\gamma\gamma} \pi^+ \pi^-)(K_S^0 \rightarrow \pi^+ \pi^-)$
 - Background
 - $B^0 \rightarrow \eta'(\rightarrow \eta_{\gamma\gamma} \pi^+ \pi^-)(K_S^0 \rightarrow \pi^0 \pi^0)$
- 3 $B^0 \rightarrow \eta'(\rightarrow \eta_{3\pi} \pi^+ \pi^-)(K_S^0 \rightarrow \pi^+ \pi^-)$
 - Background
 - $B^0 \rightarrow \eta'(\rightarrow \eta_{3\pi} \pi^+ \pi^-)(K_S^0 \rightarrow \pi^0 \pi^0)$
- 4 Summary and outlook

With some comparison to $B^0 \rightarrow \phi K^0$ (A.Gaz) and $J/\psi(\rightarrow 2\mu)K_S^0$ channels

$B^0 \rightarrow$	BR 10^{-5}	Eff. sel. %	SXF %	Bgnd.* $\cdot 10^{-6}$	Δt reso
$\eta'(\rightarrow \eta_{\gamma\gamma}\pi^+\pi^-)K_S^0$	1.1	31.9 K_S^{+-} 13.1 K_S^{00}	1.2 2.1	2.43 1.25	2.25 ps
$\eta'(\rightarrow \eta_{3\pi}\pi^+\pi^-)K_S^0$	0.6	12.5 K_S^{+-} 6.0 K_S^{00}	3.5 3.8	0.58 - [†]	1.90 ps
$\phi(\rightarrow K^+K^-)K_S^0$	0.6	35.2 K_S^{+-} 13.7 K_S^{00}		~ 20 ~ 4	2.11 ps
$\phi(\rightarrow 2\pi)K_S^0$	0.07	28.3 K_S^{+-}		~ 700	1.42 ps
$J/\psi(\rightarrow 2\mu)K_S^0$	52	-	-	-	0.92 ps

* NB. w/o continuum suppression cut!

[†]: bug found in background skimming

- Study is still very preliminary
- So far only looked at signal $\eta' \rightarrow \eta \pi^+ \pi^-$, with $\eta_{\gamma\gamma}$ and $\eta_{3\pi}$;
 - ▶ $\rho\gamma$ final state not yet addressed;
 - ▶ K_L^0 not yet looked at, too;
 - ▶ selection still to be optimized, but first attempt is good enough to start with;
- first look at continuum background (from MC5 campaign);
 - ▶ No continuum suppression yet
- need to look at peaking background;
- signal extraction and fit to be implement
 - ▶ synergies with $B^0 \rightarrow \phi K^0$ analysis;
- Still long road toward a full scale sensitivity exercise, took first steps

Additional or backup slides

- Release used `rel-00-05-03`
- code in GIT https://github.com/lacaprara/b2pd_analysis
- data used:
 - ▶ Privately produced signal;
 - ▶ MC5 for signal (BGx0) (still partially);
 - ▶ MC5 BGx1 sample for continuum background;
 - ★ Still only a fraction of produced dataset;
 - ★ Skim (with loose selection) on large dataset;
 - ★ Full selection on skimmed events;
 - ★ trying to produce a single skim for all channels, not finalized yet
 - ★ got problem (memory and crash in mixed and charged, respectively): to be investigated;

Background selection done in two steps:

- ① **Skim:** require a reconstructed decay chain, in any of the four channels, w/ loose selection and w/o vertex fit and constraint (for speed);
 - ▶ Trying to run skim on continuum background once for all channels;
- ② **Selection:** apply all selection cuts
 - 2.1 **Pre-selection:** re-reconstruct the proper decay chain (exclusive) w/ vertex and mass constraint
 - 2.2 **Final:** apply all selection cuts
 - ▶ NB No cut (yet) on continuum discriminating variable
 - ▶ still learning how to use it
 - ▶ Alessandro showed some issue in $B^0 \rightarrow \phi K^0$ presentation on 2015/12/11

$$B^0 \rightarrow \eta'(\rightarrow \eta_{\gamma\gamma} \pi^+ \pi^-)(K_S^0 \rightarrow \pi^0 \pi^0)$$


- **preselection** at least one decay chain have been reconstructed

- **good candidate selection:**


- ▶ $M_{bc} > 5.25 \text{ GeV}$;
- ▶ $\Delta E \in [-0.15, 0.25] \text{ GeV}$; (< 0.1 for $K_S^0 \rightarrow \pi^+ \pi^-$)
- ▶ $M(\eta_{\gamma\gamma}) \in [0.45, 0.57] \text{ GeV}$;
- ▶ $M(\eta') \in [0.93, 0.98] \text{ GeV}$;
- ▶ $M(\pi^0) \in [0.1, 0.15] \text{ GeV}$;
- ▶ $M(K_S^0 \rightarrow \pi^0 \pi^0) \in [0.42, 0.52] \text{ GeV}$;
- ▶ $\text{PID}_{\pi}(\pi^{\pm}) > 0.2$;
- ▶ $d_0(\pi^{\pm}) < 0.08 \text{ mm}$;
- ▶ $z_0(\pi^{\pm}) < 0.15 \text{ mm}$;
- ▶ $N_{\text{hits}_{PXD}}(\pi^{\pm}) > 1$
- ▶ $\text{P-value}_{\text{vtx}}(B_0, \eta', \eta, K_S^0) > 1 \cdot 10^{-5}$

- if $N_{\text{cands}} > 1$, select candidate with highest $\text{P-value}_{\text{vtx}}(B_0, \eta', \eta, K_S^0)$


Distributions: all/good candidates



- combinatorics larger but still manageable
- lower tails for $M_{\pi^0, K_S^0, \eta;}$
- still most of all good candidate(s) are true;





Most best candidates are true: SXF is small $\sim 2\%$



Efficiency %







preselection	38.2
good cand's	15.3
best cand	13.1

$$\epsilon(K_S^0 \rightarrow \pi^0 \pi^0) \sim 0.5 \epsilon(K_S^0 \rightarrow \pi^+ \pi^-)$$



Flavour tagging is random,
with $\epsilon(1 - 2w)^2 = 32\%$ dilution







$$B^0 \rightarrow \eta'(\eta(\gamma\gamma) \pi^+ \pi^-) K_S^0(\pi^0 \pi^0)$$



$$B^0 \rightarrow \eta'(\rightarrow \eta_{3\pi}\pi^+\pi^-)(K_S^0 \rightarrow \pi^0\pi^0)$$

- **preselection** at least one decay chain have been reconstructed
- **good candidate selection:**
 - ▶ $M_{bc} > 5.25$ GeV;
 - ▶ $\Delta E \in [-0.15, 0.25]$ GeV;
 - ▶ $M(\eta_{3\pi}) \in [0.52, 0.57]$ GeV;
 - ▶ $M(\eta') \in [0.93, 0.98]$ GeV;
 - ▶ $M(\pi^0) \in [0.1, 0.15]$ GeV;
 - ▶ $M(K_S^0 \rightarrow \pi^0\pi^0) \in [0.4, 0.52]$ GeV;
 - ▶ $\text{PID}_\pi(\pi^\pm) > 0.2$;
 - ▶ $d_0(\pi^\pm) < 0.08\text{mm}$;
 - ▶ $z_0(\pi^\pm) < 0.15\text{mm}$;
 - ▶ $N_{\text{hits}_{\text{PXD}}}(\pi^\pm) > 1$
 - ▶ $\text{P-value}_{\text{vtx}}(B_0, \eta', \eta) > 1 \cdot 10^{-5}$
- if $N_{\text{cands}} > 1$, select candidate with highest $\text{P-value}_{\text{vtx}}(B_0, \eta', \eta, K_S^0)$







$$B^0 \rightarrow \eta'(\eta(\pi^+\pi^-\pi^0) \pi^+\pi^-) K_S^0(\pi^0\pi^0)$$





- NB: channel not used in Belle
- combinatorics is huge
- η and η' reconstruction is good

Distributions: best candidate

$$B^0 \rightarrow \eta'(\eta(\pi^+\pi^-\pi^0)\pi^+\pi^-)K_S^0(\pi^0\pi^0)$$



Sizeable SXF $\sim 30\%$, smarter best cand selection is possible





Private production







Efficiency %

preselection	35.5
good cand's	10.1
best cand	5.96
SXF	3.8

$$\epsilon(K_S^0 \rightarrow \pi^0\pi^0) \sim 0.5 \epsilon(K_S^0 \rightarrow \pi^+\pi^-)$$



$$B^0 \rightarrow \eta'(\eta(\pi^+\pi^-\pi^0) \pi^+\pi^-) K_S^0(\pi^0\pi^0)$$



- [CLEO(1998)] CLEO. Observation of high momentum η' production in B decays. *PRL*, 81:1786, 1998. doi: 10.1103/PhysRevLett.81.1786. URL <http://link.aps.org/doi/10.1103/PhysRevLett.81.1786>.
- [Urquijo(2015)] Phillip Urquijo. Comparison between belle ii and lhcb physics projections. Technical Report BELLE2-NOTE-PH-2015-004, Apr 2015.
- [BABAR(2009)] BABAR. Measurement of time dependent cp asymmetry parameters in B^0 meson decays to ωK_S^0 , $\eta' K^0$, and $\pi^0 K_S^0$. *PRD*, 79:052003, 2009. doi: 10.1103/PhysRevD.79.052003. URL <http://link.aps.org/doi/10.1103/PhysRevD.79.052003>.
- [Belle(2007)] Belle. Observation of time-dependent cp violation in $B^0 \rightarrow \eta' K^0$ decays and improved measurements of cp asymmetries in $B^0 \rightarrow \varphi K^0$, $K_S^0 K_S^0 K_S^0$ and $B^0 \rightarrow j/\psi K^0$ decays. *PRL*, 98:031802, 2007. doi: 10.1103/PhysRevLett.98.031802. URL <http://link.aps.org/doi/10.1103/PhysRevLett.98.031802>.
- [Belle(2014)] Belle. Measurement of time-dependent cp violation in $b_0 \rightarrow \eta' k_0$ decays. *Journal of High Energy Physics*, 2014 (10):165, 2014. doi: 10.1007/JHEP10(2014)165. URL <http://dx.doi.org/10.1007/JHEP10%282014%29165>.