

(Second) Approval for JME-10-005

CMS MET Performance in Events Containing Electroweak Bosons from
pp Collisions at $\sqrt{s} = 7$ TeV

Jim Alexander¹, Artur Apresyan², **Stefano Lacaprara**³

¹Cornell, ²CALTECH, ³INFN Padova

JetMet meeting
CERN, 23 August 2010



Outline



- 1 Introduction
 - Basic selection
 - Pile Up
- 2 $\cancel{E}_T$ in Photon + Jet Events
 - Goal and selection
 - Results
- 3 $\cancel{E}_T$ reconstruction in events with a W boson
 - q_T uncertainties
 - $W \rightarrow e\nu$
 - $W \rightarrow \mu\nu$
- 4 Comments
- 5 Conclusion



Introduction



Goal

Demonstrate the performance of various E_T algorithms using events containing a W, Z, or high p_T photon

- The focus of this PAS is NOT to study/measure EWK bosons as such but to study MET reconstruction in those events
- Study and compare the performance of various MET algorithms in events with real MET (W), measure MET scale and resolution in events with γ/Z
- CaloMET (raw, TypeI/II corrected), TcMET, PfMET

Links:

- CADI <http://cms.cern.ch/iCMS/analysisadmin/cadi?ancode=JME-10-005>
- HN: <https://hypernews.cern.ch/HyperNews/CMS/get/JME-10-005.html>
- Twiki: <https://twiki.cern.ch/twiki/bin/view/CMS/EwkMetComm>
- Previous Approval (indico) - Pre-Approval (indico)



History



- pre-approval (with $\sim 12 \text{ nb}^{-1}$) on June 28
- approval on July 9th with $\sim 56 \text{ nb}^{-1}$ **NOT approved**
 - ▶ While one has to congratulate all people involved for the fast production of the plots with the newly arrived data, we also saw that **many questions have come up and issues need to be understood in the plots, which were not visible before.**
 - ▶ more work has to go into the understanding of the new results, **in particular for the gamma+jet sample.**
 - ▶ ... **Concentrate on the data which have been taken up to now.**
 - ▶ ... it is clear that **we start to see PU effects**, thus certain plots should be done as a function of Nvtx ...
- Decision to skip ICHEP and concentrate on analysis for $\gamma + \text{jet}$ and PU issues.
- **Second pre-approval on August, 9th**
- **Timescale for this 2nd approval aims at PIC2010 (1-4 Sept)**



A Lighter PAS



Statistics

- Stay with “pre-ICHEP” integrated luminosity, as suggested.
 $\int \mathcal{L} dt \sim 200 \text{ nb}^{-1}$.

New PAS

- $\cancel{E}_T$ in Photon + Jet Events
- $\cancel{E}_T$ reconstruction in events with a W
- POSTPONED WAITING FOR MORE DATA Performance of $\cancel{E}_T$ reconstruction in events with a Z boson
- DROP FOR THE TIME BEING
 - ▶ Effects of muon reconstruction uncertainties on $\cancel{E}_T$
 - ▶ $\cancel{E}_T$ significance
 - ▶ Estimating the $\cancel{E}_T$ distribution in $W \rightarrow e\nu$ events



Title and abstract



CMS MET Performance in Events Containing Electroweak Bosons from pp Collisions at $\sqrt{s} = 7$ TeV

JME-10-005

During the spring of 2010, the LHC delivered proton-proton collisions with a centre-of-mass energy of 7 TeV. In this note, we present results of studies of missing transverse energy, as measured by the CMS detector, in events containing W bosons or isolated, high transverse momentum photons. The performance of several different MET reconstruction algorithms is compared.

Editors: Artur Apresyan, Stefano Lacaprara, Jim Alexander (senior)
ARC: Fabio Cossutti (Trieste), Sharon Lee Hagopian (Florida-state),
Paraskevas Sphicas [chair] (CERN)



Supporting Documents

Several supporting AN

- **AN-2010/118** CMS MET Performance in Events Containing Electroweak Bosons decaying into muons from pp Collisions at $\sqrt{s} = 7$ TeV (Padova)
- **AN-2010/131** Type-I and Type-II CaloMET performances in 7TeV data (Hamburg, TTU, Florida)
- **AN-2010/132** MET Scale Validation with Photon + Jet Events (Texas Tech Uni.)
- **AN-2010/176** Commissioning of the missing transverse energy in $W \rightarrow \mu\nu$ events for 12 nb^{-1} with the pp center-of-mass energy of $\sqrt{s} = 7$ TeV (Brown)
- **AN-2010/202** Missing transverse energy performances with electroweak bosons decaying into electrons in pp collisions at $\sqrt{s} = 7$ TeV (Saclay)



Changes since second Pre-approval

- Add Pile-up vs no Pile-up plots for key distribution (γ +jet response and resolution)
- Add number of events selected for each analysis
- Update to 36x simulation for $tc\cancel{E}_T$
- State more clearly that we use Calo $\cancel{E}_T$ type II only.
- Minor fix in plots style (no change in content)
- ...

Comments recieved

Will report major comment and our reply near the end of presentation



Basic Selection and Definition

Uniform Selection

- Vertex requirement, datasets, trigger selections (muon and electron)
- Electron and muon IDs following the VBTF recommendations (more on this later...)
- EGamma electrons are used in the studies of PFMET, in agreement with PF POG
- ECAL/HCAL noise is cleaned in re-reco used in the analysis




Figure: Kinematics: (a) Photon-Jet events; (b) W events.




○○○○○○○○○○

○○○○○○○○○○○○○○

PU Estimation and Treatments



- MET related variables are sensitive to PU.
- **Require just one Primary Vertex.**
- MOTIVATION: start with simple (no PU) events and then eventually move to events with PU (not for this PAS)
- Estimate PU distribution by PV multiplicity: **58% 1 PV**




- Contamination from 2 not resolved PV estimated with toy MC (next slide)



Contamination from multiple PV



- Get z_{PV} distribution from data: $\sigma = 6.25 \pm 0.26$ cm



- Get minimal Δz for two PV to be resolved from data:
 $\Delta z = 1.5 \pm 0.5$ cm
- Use Toy MC to estimate the $PV = 2$ contamination in $PV = 1$ sample: $5 - 7 \pm 2\%$. Negligible for $PV > 2$.
- scale the multi-PV distribution and subtract from the 1-vertex distribution



Outline

- 1 Introduction
 - Basic selection
 - Pile Up
- 2 $\cancel{E}_T$ in Photon + Jet Events
 - Goal and selection
 - Results
- 3 $\cancel{E}_T$ reconstruction in events with a W boson
 - q_T uncertainties
 - $W \rightarrow e\nu$
 - $W \rightarrow \mu\nu$
- 4 Comments
- 5 Conclusion



$\cancel{E}_T$ in Photon + Jet Events



Goal

- Events with no intrinsic $\cancel{E}_T$
 - cross-section larger than that of Z
 - *induce* $\cancel{E}_T$ by removing γ deposit in calo
 - magnitude of *induced* $\cancel{E}_T$ well known.
-
- trigger HLT_PHOTON10_L1R
 - Photon ID based on *loose* selections
(see ‘‘Photon reconstruction and identification at $\sqrt{s} = 7\text{ TeV}$ ’’, EGM-10-005.)

Analyzed Data Sample:

$$\int \mathcal{L} dt = 198.1 \text{ nb}^{-1}$$


Event selection:

- ECAL energy deposit ($\Delta R < 0.4$) $E < 4.2 + 0.004 \times q_T$.
- HCAL energy deposit ($\Delta R < 0.4$) $E < 2.2 + 0.001 \times q_T$.
- Ratio HCAL/ECAL ($\Delta R < 0.15$) $R < 0.05$.
- N. tracks ($\Delta R < 0.4$) $N_{trk} < 3$.
- $\Sigma_{(0.04 < \Delta R < 0.4)} p_T < 2.0 \text{ GeV} + 0.001 \times q_T$
- $R9 > 0.9 \times E^\gamma$
- γ cluster major and minor 2^{nd} moments in $0.20 - 0.35$, $0.15 - 0.3$.
- $\eta_{width} < 0.03$
- $q_T > 20$ and $|\eta| < 1.479$ (Barrel)
- Only 1 Primary Vertex (No PU)
- γ supercluster not match pixel hits consistent with a track from the interaction region ($W \rightarrow e\nu$ suppression)




Photon q_T and $W \rightarrow e\nu$ suppression

CMS preliminary 2010

 $\sqrt{s} = 7$ TeV

- significant amount of $W \rightarrow e\nu$ events pass standard γ loose ID
- not considered as background in previous approval
- strongly suppressed ($\sim 98\%$) by pixel seeds veto

CMS preliminary 2010

 $\sqrt{s} = 7$ TeV




q_T distribution of events selected as photon-jet candidates. Monte Carlo predicted rates for signal and backgrounds are as shown.

QCD di-jet contamination is dominated by $\pi^0 \rightarrow \gamma\gamma$ enriched jets. Still good for energy scale studies



Effect of di-jet contamination (NOT FOR PAS)

Shown response ($u_{||}/q_T$) for MC pure $\gamma + jet$ and $\gamma + jet$ & QCD di-jet



MC studies shows that QCD (di-jet) contamination give no bias in MET response



Recoil projections along the γ axis

Parallel component used to study $\cancel{E}_T$ scale and resolution **PAS**





Figure: Decomposition of hadronic recoil into components parallel and perpendicular to photon probe. Upper row: $u_{||}$ distributions for (left to right) $\text{calo}\cancel{E}_T$, $\text{tc}\cancel{E}_T$, and $\text{pf}\cancel{E}_T$;

$W \rightarrow e\nu$ contamination strongly suppressed by pixel seed veto



Recoil projections perpendicular to the γ axis


Perpendicular component used to study $\cancel{E}_T$ resolution due to calo noise, UE and PU **PAS**



- PU affect the width of distribution See later
- 1 Primary vertex to select event with no PU
- NO PileUp in MC simulation




Angular correlation between $\vec{u}_T$ and q_T



- $W \rightarrow e\nu$ is flat in $\Delta\phi$: large contribution at low $\Delta\phi$ if not suppressed (see last approval talk).
- Residual discrepancies at low $\Delta\phi$ is likely to be a resolution effect in the tail of the distribution;
- Not present in $p\cancel{E}_T$;
- few % or few ‰ effect;
- Does not affect results on response or resolution




E_T Scale: $|u_{||}|/q_T$



- JES correction for calo E_T based on q/g jets. Response to q jets is $\sim 10\%$ higher than g one. Direct γ have mostly q jet, so overcorrection is expected;
- Tc E_T and pf E_T are not corrected for JES, since the effects, already in the algo;
- also shown ≥ 2 PV distribution: response is not sensitive to Pile-UP **NEW**




Resolution (RMS) for $u_{\perp}$ vs q_T



- including uncertainties (statistical) from MC
- Resolution corrected for response curve
- also shown ratio $\geq 2PV / 1PV$: resolution IS sensitive to PU **NEW**
- Interesting to note that the PU effect is $\sim$ the same for all algo.
- Also $\sim$ independent of q_T
- Measuring $\cancel{E}_T$ resolution in the data



Resolution (RMS) for $u_{||}$ vs q_T



- including uncertainties (statistical) from MC
- Resolution corrected for response curve
 - ▶ measure RMS of $(u_{||}/q_T)$ and then multiply by average bin q_T and scale correction
- also shown ratio $\geq 2PV / 1PV$: resolution IS sensitive to PU **NEW**
- Use of tracking information improves significantly the MET resolution



Outline



- 1 Introduction
 - Basic selection
 - Pile Up
- 2 $\cancel{E}_T$ in Photon + Jet Events
 - Goal and selection
 - Results
- 3 $\cancel{E}_T$ reconstruction in events with a W boson
 - q_T uncertainties
 - $W \rightarrow e\nu$
 - $W \rightarrow \mu\nu$
- 4 Comments
- 5 Conclusion





W q_T uncertainty



MC uncertainties for q_T

- Use standard PYTHIA and POWHEG
- Compare the two q_T distribution
- Use difference between original and reweighted as systematic error, bin per bin.
- add this to error from PU contamination, as described before






$W \rightarrow e\nu$ selection



- Standard VBTF selection WP 80% no $\Delta\eta$ cut in endcap (EGamma prescription)
 - ▶ HLT $p_T(e) > 10$ GeV
 - ▶ Electron id 80% efficiency.
 - ▶ $|\eta_e| < 2.5$ excluding $1.4442 < |\eta| < 1.56$
 - ▶ η dependent isolation on ECAL, HCAL and tracksAN-2010/133
 - ▶ No second electron $p_T > 20$ GeV
- GSF filter + supercluster $p_T > 25$ GeV (VTBF is > 20)
- only 1 Primary vertex
- PU contamination cleaning
- Additional cuts to enrich $W \rightarrow e\nu$
 - ▶ $\cancel{E}_T > 25$ GeV
 - ▶ $M_T > 50$ GeV
- POWHEG MC used
- QCD and EWK normalization by a fit on $\cancel{E}_T$ shape
- $\int \mathcal{L} dt = 255 \text{ nb}^{-1}$, 461 events selected



$\cancel{E}_T$ distribution in $W \rightarrow e\nu$ events




REMINDER: changes from previous PAS:

- $p_T(ele) > 25 \text{ GeV}$ (was 20)
- NO $\cancel{E}_T$ cut;
- 1 primary vertex and PU cleaning
- use this distribution to normalize QCD and EWK in MC.




Recoil in $W \rightarrow e\nu$ events



- agreement at low $u_{\perp}$ much better due to Primary Vertex requirement and PU cleaning
- Uncertainties at low $u_{\perp}$ dominated by q_T ones.




Recoil along q_T in $W \rightarrow e\nu$ events



- Projection along $p_T(\ell)$, not q_T (unknown): correlation is good for boosted W
- Asymmetry due to strict isolation cut on $W \rightarrow e\nu$: When $u_{\parallel}$ is positive, electron and hadronic activities are in the same hemisphere, more likely that the electron is not isolated.
- Tail at low $u_{\parallel}$: $W \rightarrow e\nu$ events with boosted W .




Recoil perpendicular to q_T in $W \rightarrow e\nu$ events



- Calo $\cancel{E}_T$ is clearly broader than TC and PF $\cancel{E}_T$;
- Still dominated by true $\cancel{E}_T$, width similar to that of $u_{\parallel}$.



$\Delta\phi$ recoil-lepton in $W \rightarrow e\nu$ events



- Good agreement within statistical errors



$W \rightarrow \mu\nu$ selection



- **Standard VBTF selection**

- ▶ $\text{HLT } p_T(\mu) > 9 \text{ GeV}$ HLT_MU9
- ▶ Muon *Global* and *Tracker*
- ▶ Tracker hits > 10, Pixel hits > 0; Muon hits > 0;
- ▶ EM veto < 4 GeV; Hadronic veto < 6 GeV;
- ▶ Relative combine isolation < 0.15
- ▶ impact parameter (beam spot) < 2 mm
- ▶ Global fit $\chi^2 < 10$
- ▶ $|\eta| < 2.1$

- **Muon $p_T > 25$**

- **only 1 Primary vertex**

- PU contamination cleaning

- Additional cuts to enrich $W \rightarrow \mu\nu$


- ▶ $\cancel{E}_T > 25 \text{ GeV}$
- ▶ $M_T > 50 \text{ GeV}$

- **QCD and EWK normalization by a fit on $\cancel{E}_T$ shape**

- **$\int \mathcal{L} dt = 246 \text{ nb}^{-1}$ 514 $W \rightarrow \mu\nu$ selected**



$\cancel{E}_T$ distribution in $W \rightarrow \mu\nu$ events




REMINDER: changes from previous PAS:

- $p_T(\mu) > 25$ GeV (was 20)
- NO $\cancel{E}_T$ cut;
- 1 primary vertex and PU cleaning
- use this distribution to normalize QCD and EWK in MC.



Recoil in $W \rightarrow \mu\nu$ events




- Good agreement at low u_T thanks to PU contamination removal;

○○○○
○○○○○○○○○○

○○○○○○○○○○●○○




Recoil along q_T in $W \rightarrow \mu\nu$ events



- Small/no asymmetries as compared to $W \rightarrow e\nu$ due to softer isolation cut.
- As for $W \rightarrow e\nu$, good correlation between $p_T(\ell)$ and q_T only for boosted W




Recoil perpendicular to q_T in $W \rightarrow \mu\nu$ events



- Narrower distribution for TC and PF $\cancel{E}_T$



$\Delta\phi$ recoil-lepton in $W \rightarrow \mu\nu$ events



- Good agreement between data and MC



Outline



- 1 Introduction
 - Basic selection
 - Pile Up
- 2 $\cancel{E}_T$ in Photon + Jet Events
 - Goal and selection
 - Results
- 3 $\cancel{E}_T$ reconstruction in events with a W boson
 - q_T uncertainties
 - $W \rightarrow e\nu$
 - $W \rightarrow \mu\nu$
- 4 Comments
- 5 Conclusion



Comments #1

Which is the size of PU effect??

The current PAS tends to leave the reader hanging whether there is or is not an understanding of MET for EWK events with pile-up at the level of the ICHEP data sample since it explicitly only addresses events with 1PV (see statement on lines 20-23) ... Nevertheless, I would propose that it is still beneficial to include the results from data alone so the reader can judge the size of the effect rather than not include anything.


Added plot comparing data w/ and w/o PU for γ +jet.
Not for W plots: overkill and statistically not very significant.



Comments #2

this analysis measures the response and resolution of the hadronic recoil. I think to call it a missing ET response measurement is a bit misleading because it doesn't transfer to other event topologies with missing ET

True: results does not work for $\cancel{E}_T$ in **all** event topologies, but are fine when $\cancel{E}_T$ is dominated by hadronic topologies. Keep the name. Example are Z events, not included in this PAS but will do in future one. Already shown that responses for $\gamma + jet$ and Z compared very well .





Comments #3

 $\cancel{E}_T$ response misleading

It seems to me the resolution should be measured from $\text{RMS}(u_{\parallel}-q_T)$ because the width of the $u_{\parallel}$ spectrum may be driven by the width of the photon q_T bin rather than the recoil resolution

The resolutions of $u_{\parallel}$ are calculated using $(u_{\parallel}/q_T)$, binned in q_T . For a given bin we measure RMS of $(u_{\parallel}/q_T)$ and then multiply this quantity by (average bin q_T) $\times$ (scale correction).

This allows to measure the MET resolutions corrected for the scale, measuring the 2^{nd} central moment, that is simply shifted to 1.0 instead of being at 0.



Comments #4



W Analysis Scope

The main point of the W analysis is to demonstrate the agreement of data and MC simulation. I do not think that you can measure the recoil response nor the resolution using W decays because you have nothing to calibrate the recoil against.

True. This section of the current PAS focuses on MET reconstruction in W events not response or resolution. Our W plots are mainly to demonstrate a good data/MC comparison and a simple comparison of performance of MET algos.

The response and resolution measurements using Z events were in the initial plan for this PAS, but due to the limited statistics, were dropped.



Comments #5



I think it is not clear that the widths of $u_{\parallel}$ and $u_{\perp}$ are dominated by the recoil resolution. Both are projections of the recoil p_T on an axis that is only weakly correlated with the direction of the recoil p_T . The range of $u_{\parallel}$ and $u_{\perp}$ is $-\max(u_T)$ to $+\max(u_T)$. So the recoil p_T distribution will affect the width of the $u_{\parallel}$ and $u_{\perp}$ distributions.

The RMS of either $u_{\parallel}$ or $u_{\perp}$ will in general contain two pieces: a narrow core reflecting the basic hadronic resolution, and a wide (and in practice slightly asymmetric) component that includes the contamination from actual p_T . (this second component is $\in [-\max(u_T), +\max(u_T)]$) It is true that we haven't really probed this issue very deeply, and consequently the text is carefully written to offer only a qualitative, not quantitative, interpretation of the plots. Nevertheless we feel the text offers some insight into the physical content of the plots, and for that reason should stay



Outline



- 1 Introduction
 - Basic selection
 - Pile Up
- 2 $\cancel{E}_T$ in Photon + Jet Events
 - Goal and selection
 - Results
- 3 $\cancel{E}_T$ reconstruction in events with a W boson
 - q_T uncertainties
 - $W \rightarrow e\nu$
 - $W \rightarrow \mu\nu$
- 4 Comments
- 5 Conclusion



Conclusion

$\gamma + jet$

- Studies presented with 198.1 nb^{-1}
- Analysis updated
 - ▶ considering $W \rightarrow e\nu$ contamination
 - ▶ introducing further cut (pixel seeds) to effectively reduce it
 - ▶ reducing PU effect by requiring just 1 Primary Vertex and multi-PV cleaning
- Data driven assessment of $\cancel{E}_T$ scale and resolution.
- PU effect visible in MET resolution but NOT in response **NEW**
- Good agreement data-MC seen



Conclusion/II

$W \rightarrow e\nu$ and $W \rightarrow \mu\nu$

- Studies presented with 255 and 246 nb⁻¹ respectively
- Improvement wrt previous PAS
 - ▶ stricter cuts to select purer W sample
 - ▶ Select events with just one Primary Vertex
 - ▶ clean PU contamination in 1-vertex sample
- Uncertainties included:
 - ▶ q_T spectra using PYTHIA and POWHEG
 - ▶ from PU contamination
- Good agreement data-MC seen



Conclusion/III



PAS conclusion

- The performance of three $\cancel{E}_T$ algorithms have been examined with early data
- emphasis on the calibration scale and resolution of the $\cancel{E}_T$ response.
- Very good agreement between data and MC
- the improvement that results from the inclusion of charged particle tracking in jet reconstruction is visible and significant.
- The difference in performance is further confirmed in $\cancel{E}_T$ distributions of $W \rightarrow \ell \nu$ event samples which contain genuine $\cancel{E}_T$.



BACKUP




Angular correlation γ jet: big bins

Typell Calo $\#_T$, TC $\#_T$, PF $\#_T$


CMS preliminary 2010

$\sqrt{s} = 7$ TeV




CMS preliminary 2010

$\sqrt{s} = 7$ TeV






CMS preliminary 2010

$\sqrt{s} = 7$ TeV






Angular correlation γ jet: statistics




small/no statistics at low $\Delta\phi$ for low $\hat{p}_T$ bins for QCD di-jet events



$\gamma + \text{jet}$ angular correlation with $E_\gamma > 30$ GeV



After the “pixel seeds” cleaning. $PT > 30$ cut is applied to enrich the sample with uncleaned W. Main W contamination is expected at $30 < PT < 45$. Instead somewhat better DATA/MC agreement






TC#_T: $\gamma + jet$ angular correlation at different q_T



Left: $20 < q_T < 50$ GeV, Center: $35 < q_T < 100$ GeV, Right: $50 < q_T < 100$ GeV



Possible NLO vs LO effect at low q_T ?



CaloMET: $\gamma + jet$ angular correlation at different q_T



Left: $20 < q_T < 50$ GeV, Center: $35 < q_T < 100$ GeV, Right: $50 < q_T < 100$ GeV

CMS preliminary 2010




$\sqrt{s} = 7$ TeV

CMS preliminary 2010

$\sqrt{s} = 7$ TeV

CMS preliminary 2010







$\sqrt{s} = 7$ TeV





Any effect of PU in EWK studies?



- Very limited effect of PU on M_T and $\cancel{E}_T$ distribution, on which the EWK analysis is based. Significant only for hadronic recoil
- Below: $\cancel{E}_T$ (l), M_T (c) and u_T (r) for $W \rightarrow \mu\nu$ case with $\text{Pf}\cancel{E}_T$





Any effect of PU in EWK studies? /II


- From $\gamma + jet$ analysis, no (or negligible) effect on MET scale from the PU.
- (limited) effect on $\cancel{E}_T$ resolution from PU
- will be needed for analysis such as W mass measurement (not in ICHEP EWK program)






Effect of PU on resolution

Resolution (RMS for $u_{\perp}$ and $u_{\parallel}$) as a function of q_T for events with 1 Primary Vertex and ≥ 2 PV for three different $\cancel{E}_T$ algos. Not for PAS Data after $W \rightarrow e\nu$ pixel veto






Effect of pixel cleaning on Resolution




Before the "pixel seeds" cleaning

caloMET Typell






tcMET







pfMET





After the "pixel seeds" cleaning







Effect of QCD contamination on Resolution










$u_{||}$ Barrel vs Endcap for TcMet








Effect of PU contamination on Recoil $W \rightarrow e\nu$

Pileup effects on caloT2 recoil variables (data only)





Get normalized shapes






Any effect of PU in EWK studies? $W \rightarrow e\nu$









Any effect of PU in EWK studies? $W \rightarrow \mu\nu$







MT distribution for $W \rightarrow \mu\nu$





Muon p_T distribution for $W \rightarrow \mu\nu$





For 20 and 25 p_T cut

Note: different QCD/EWK normalization for MC from MET distribution



Recoil perpendicular to q_T in $W \rightarrow \mu\nu$ events







Comparison with EWK PAS at ICHEP $M_T W \rightarrow \mu\nu$



Left EWK PAS , Right Current analysis





- $p_T > 25 \text{ GeV}$ (VTBF is > 20)
- $\cancel{E}_T > 25 \text{ GeV}$ (VTBF no cut)
- only 1 Primary vertex
- Shown only for $M_T > 50$
- Plot NOT for current PAS!



Comparison with EWK PAS at ICHEP $\cancel{E}_T W \rightarrow e\nu$



Left EWK PAS $\cancel{E}_T$, Right current work $\cancel{E}_T$



- $p_T > 25 \text{ GeV}$ (VTBF is > 20)
- only 1 Primary vertex