# The Magnetic field of the Earth and Planets

# Earth's magnetic field

#### 1. Structure

Earth's magnetic field has many relevant effects on the space around the planet. It protects the surface from the solar wind creating the *magnetosphere*, it traps charged particles in the *Van Allen radiation belts* and it is involved in the formation of the spectacular *aurora borealis* (or northern lights). In Fig. 1 the intensity of the field is shown at different locations on the Earth surface.



Fig. 1: magnetic field of the Earth according to IGRF (International Geomagnetic Reference Field, year 2000). Red color corresponds to 68000 nT while blue color is for 24000 nT.

The IGRF representation of the magnetic field outside the Earth is obtained by a spherical harmonics development. Starting from the Ampere-Maxwell equation:

$$\nabla \times \overline{B} = \mu_0 \left( j + \varepsilon_0 \frac{\partial \overline{E}}{\partial t} \right)$$

we can assume that outside the Earth surface j=0 (the source is inside the Earth). We can also adopt the approximation that  $\left(\frac{\partial \overline{E}}{\partial t}\right) = 0$  so that we get:

$$\nabla \times \overline{B} = 0$$

As a consequence, the magnetic field can be derived from the gradient of a scalar function V that we call magnetic potential:

$$\overline{B} = -\nabla V$$

Moreover, we know that the third Maxwell equations requires that:

$$\nabla \cdot \overline{B} = 0$$

which leads to a Laplace equation for the magnet field potential V out of the Earth's surface:

 $\nabla^2 V = 0$ 

Due to the spherical nature of the boundary conditions the solution can be expressed a series of spherical harmonics:

$$V(r,\theta,\varphi) = a \sum_{n=1}^{k} \left(\frac{a}{r}\right)^{n+1} \sum_{m=0}^{n} \left(g_{n}^{m} \cos m\varphi + h_{m}^{n} \sin m\varphi\right) P_{n}^{m}(\theta)$$

where *a* is the Earth's radius (for the *International Geomagnetic Reference Field* (IGRF) a=6371,2 km);  $g_n^m e h_n^m$  are the Gauss' coefficients and r,  $\theta$ ,  $\varphi$  are the radial distance, co-latitude (complementary angle of the latitude) and longitude measured from Greenwich towards east, respectively. The  $P_n^m$  are the Legendre functions. The Gauss' coefficients are given by IGRF and are computed with least squares interpolation of measured data on the magnetic field. The components of the magnetic field are given by:

$$B_{r} = \frac{-\partial V}{\partial r}$$
$$B_{r} = \frac{-1}{r} \frac{\partial V}{\partial \theta}$$
$$B_{\phi} = \frac{-1}{r \sin \theta} \frac{\partial V}{\partial \phi}$$

#### 2. Source

What is the source of the magnetic field? To answer this question we have to explore the interior of the Earth. Earthquakes and the difference in the propagation of seismic waves which depends on the composition of the crossed material, have lead to the following model of Earth's interior:



Fig. 2: different propagation paths of seismic waves on the interior of the Earth. Reflection and refraction phenomena.

The Earth is divided in the lithosphere (crust and uppermost mantle), the mantle, an outer liquid core and an inner solid core.



10 wt%) with very small amounts of some lighter elements. It is the outcome of the Earth differentiation with the heavier elements sinking towards the center of the planet after its formation. It is unattached to the mantle, suspended in the molten outer core and, in spite of its high temperature (5000-6000 C), it is believed to have solified as a result of *pressure-freezing* which occurs to most liquids under extreme pressure.



Fig. 4. Typical pressure-temperature diagram for a solid (the dotted line shows the anomalous behavior of water). Increasing the pressure leads to crystallization at higher temperatures. On the left, the phase diagram for a Fe-10wt%Ni alloy from Lin et al., Geophysical Research Letters 29, 1471, 2002.

The inner core has a density of about 15 g/cm<sup>3</sup>.

*The outer core* has a composition similar to the inner core with about 10% of sulphur, oxygen and other light elements. The temperature is 4000-5000 C and the density is between 10 and 12 g/cm<sup>3</sup>. Since the pressure is lower and it is hot, the outer core is liquid. Over geologic time, the inner core grows at the expense of the outer core as the whole Earth cools. The pressure is the same but the temperature drops and the pressure-freezing radius extends farther out. In the outer core the liquid metal is electrically conducting. This conductive layer combines with *Earth's rotation* to create a

dynamo effect that maintains a system of electrical currents that creates the Earth's magnetic field. In conclusion, the Earth's magnetism is due to a combination of rotation and molten metallic core.

#### **3.** Planetary rotations and magnetic fields.

The rotation rate of a planet and the presence of a molten core are fundamental requirements for the presence of a magnetic field. Here below we list the planets of th solar system and the parameters related to rotation and dipole axis inclination. The angle  $\varepsilon$  is the obliquity, the inclination of the rotation axis of the planet respect to the orbital plane, the angle  $\alpha$  is instead the angle between the magnetic field axis (in dipole approximation) and the planet rotation axis. The rotation period is given in days for the terrestrial planets and hours for the outer planets while the magnetic field must be attributed to its slow rotation while for mars it is due to the absence of a molten core. For the outer planets the magnetic fiels has a more complex source that for Jupiter and Saturn may be related to the formation of metallic hydrogen close to the planet core.



|         | $\alpha$ (deg) | $\epsilon$ (deg) | P (days)    | M <sub>B</sub> (in Earth's<br>units) |
|---------|----------------|------------------|-------------|--------------------------------------|
| Mercury | 10             | 0                | 58.6        | 4 10-4                               |
| Venus   | 0              | 177              | -243        | 0                                    |
| Earth   | 10.8           | 23.5             | 1           | 1                                    |
| Mars    | 0              | 25.9             | 1           | 0                                    |
| Jupiter | 9.6            | 3.12             | 9.9 (hrs)   | $2  10^4$                            |
| Saturn  | < 1            | 26.75            | 10.7 (hrs)  | 600                                  |
| Uranus  | 60             | 97.86            | -17.2 (hrs) | 50                                   |
| Neptune | 47             | 29.56            | 16.1 (hrs)  | 25                                   |

# 4 Dipole approximation

The Earht's magnetic field is due by 97-99% to electric currents in the nucleus, 1-2% to magnetized rocks in the crust, and 1-2% to electric currents around the Earth. It has a strongly dipole-dominated structure (up to 90% of the field strength) with the magnetic south close to the geographic north (see figure).



Glatzmaier-Roberts numerically solved the magnetohydrodynamics equations (MHD) to model the magnetic field generation in a fluid outer core surrounding a solid inner core (Glatzmaier and Roberts, Phys. Earth Planet. Inter., 91, 63-75, 1995). They were also able to reproduce the dipole *field reversal* which occurs randomly every about 250000 years. They find that the intensity of the magnetic dipole moment decreases by about a factor of ten during the reversal and recovered immediately after, similar to what is seen in the Earth's paleomagnetic reversal record.



Fig. 5. Timing of magnetic reversals from magnetized rocks. Dark areas denote periods where the polarity matches today's polarity, light areas denote periods where that polarity is reversed.

On the basis of their model they explain how convection in the fluid outer core is continually trying to reverse the field but that the solid inner core inhibits magnetic reversals because the field in the inner core can only change on the much longer time scale of diffusion. Only once in many attempts is a reversal successful, which is probably the reason why the times between reversals of the Earth's field are long and randomly distributed (Glatzmaier and Roberts, Nature, 377, 203-209, 1995).



Fig. 6: 3D magnetic field structure simulated with the Glatzmaier-Roberts (1995) geodynamo model.

Magnetic field lines are blue where the field is directed inward (North) and yellow where directed outward (South). The rotation axis of the model Earth is vertical and through the center. The field lines are drawn out to two Earth radii.

#### 4. Magnetic dipole

The equations for a 3D magnetic dipole are:

$$B_r = \frac{2\mu_0 m}{4\pi r^3} \cos\theta$$
$$B_r = \frac{-\mu_0 m}{4\pi r^3} \sin\theta$$
$$B_{\pm} = 0$$

where *m* is the magnetic moment that, for a coil of area *As* passed through a current *i*, is defined as:

 $m = As \cdot i$ 



When dealing with the Earth's magnetic field, usually the co-latitude  $\theta$  is substituted by the latitude and the direction of the dipole moment is directed downwards. As a consequence, the formula for a dipolar magnetic field changes to:

$$B_r = \frac{-2M_B}{r^3} \sin \theta$$
$$B_\theta = \frac{M_B}{r^3} \cos \theta$$
$$B_\phi = 0$$

where  $M_B = \mu \frac{m}{4} \pi = 7.906 \cdot 10^{25} gauss/cm^3$  is the magnetic moment of the Earth. The filed lines can be derived with a simple reverse reasoning. We assume that the equation for the filed line is already known and it is  $r = r_e \cos(\theta)^2$  and we verify that the magnetic field **B** is tangent to this line. The tangent vector to the line defined by  $r = r_e \cos(\theta)^2$ , with  $r_e a$  constant, is given by:

$$\mathbf{v} = \dot{r} \, \mathbf{e}_r + r \, \dot{\theta} \, \mathbf{e}_{\theta} + r \sin \theta \, \dot{\phi} \, \mathbf{e}_{\phi} = -2 \, r_e \cos \theta \sin \theta \, \dot{\theta} \, \mathbf{e}_r + r_e \cos^2 \dot{\theta} \, \mathbf{e}_{\theta} = r_e \cos \theta \, \dot{\theta} \left(-2 \sin \theta \, \mathbf{e}_r + \cos \theta \, \mathbf{e}_{\theta}\right)$$

in the final right end side of the equation, the vector within the round brackets is parallel to the magnetic field  $\mathbf{B}$ . This means that indeed the equation for the field line is correct. We can at this point compute the value of the magnetic field along a field line as:

$$|\mathbf{B}| = \frac{M_B}{r_e^3} \cos \theta^6 \cdot [4\sin \theta^2 + \cos \theta^2]^{\frac{1}{2}} = \frac{M_B}{r_e^3} \frac{[4 - 3\cos \theta^2]^{\frac{1}{2}}}{\cos} \theta^6 = B_e \frac{\sqrt{4 - 3\cos \theta^2}}{\cos} \theta^6$$
  
where  $B_e = \frac{M_B}{r_e^3}$ .

# Motion of charged particles in the magnetic field of the Earth

The motion of charged particles trapped in the magnetic field of the Earth is multi-periodical and it can be studied with a perturbative approach. It can be divided in three different component: the gyromagnetic, drift and mirror motion. This separation is possible since the different types of motion have different timescales. The gyromagnetic motion can be considered as the unperturbed motion of particles while the drift and the mirror motion are slow perturbations of the gyromotion.

#### 1. The gyromagnetic motion

The equation of motion of a charge particle q under the action of a constant magnetic field directed along the z-axis are derived from the Lorentz's equation of the force:

$$\begin{array}{l} m \dot{v}_x = q \, v_y B \\ m \dot{v}_y = q \, v_x B \end{array} \quad \text{where} \quad \boldsymbol{B} = (0, 0, B) \quad \text{and} \quad \boldsymbol{V} = (v_x, v_y, 0) \end{array}$$

to solve this equations we can derive the equation of motion once:

$$\ddot{x} = q \frac{B}{m} \ddot{y} = -\left(\frac{qB}{m}\right)^2 \dot{x}$$
$$\ddot{y} = q \frac{B}{m} \ddot{x} = -\left(\frac{qB}{m}\right)^2 \dot{y}$$

Integrating back these equations we get:

$$\ddot{x} = -\Omega_c^2 x$$
$$\ddot{y} = -\Omega_c^2 y$$

They are the equations of uniform circular motion with frequency:

$$\Omega_c = \frac{qB}{m}$$

called cyclotron frequency. The centripetal acceleration and the radius are given by:

$$|a| = \Omega_c^2 |r| = \frac{v^2}{r}$$
 and  $r_c = \frac{v}{\Omega_c} = \frac{v_\perp m}{q B}$ 

where  $r_c$  is the radius of the circular trajectory while  $v \perp = v_x^2 + v_y^2$  is the modulus of the circular velocity perpendicular to the magnetic field direction. Ad example, electrons moving along the magnetic filed lines of the Earth have an average radius of 100 m, a period of the order of the  $\mu$ s and a velocity around 10<sup>6</sup> km/s.

#### 2. The drift motion

The drift motion of charged particles in the Earth's magnetic field is due to the curvature and gradient of the field and, to a lower extent, to the presence of the gravity field and a weak electric field. We list here below the contributions to the drift motion:

1) If the magnetic field lines are curved then:

$$v_{DI} = m v_{\parallel}^2 \frac{\boldsymbol{B} \times \boldsymbol{n}}{R_c q B^2}$$

where  $v_{\parallel}$  is the component of the particle velocity parallel to the tangent of the local magnetic field line and R\_c is the curvature radius of the field line.

2) If there is a non-zero field gradient, then:

$$v_{D2} = \frac{1}{2} m v_{\perp}^2 \frac{\boldsymbol{B} \times \boldsymbol{\nabla} \boldsymbol{B}}{q} q \boldsymbol{B}^3$$

The sign of both  $v_{D1}$  and  $v_{D2}$  depend on the charge q. As a consequence, protons and electrons drift towards different directions. Electrons drift towards west while protons towards east.

3) If there is an electric field **E** and a gravity field **G** there is a third drift velocity given by:

$$\boldsymbol{v}_{D3} = \frac{\boldsymbol{E} \times \boldsymbol{B}}{\boldsymbol{B}^2} + \frac{\boldsymbol{m}}{\boldsymbol{q} \boldsymbol{B}^2} \boldsymbol{G} \times \boldsymbol{B}$$

The dominant velocity component is vD1 which is prop velocity components of drifting charges.

#### 3. The adiabatic invariant and the mirror motio

The mirror motion can be understood in a naïve way by inspecting the side figure. When the field lines are converging and the  $v_{\parallel}$  is directed towards the closing lines, the Lorentz force related to the gyromagnetic motion presents a component which acts like a repelling force.

Within a perturbative approach, we can interpret the mirror motion as the consequence of the conservation of an adiabatic invariant, the magnetic



moment of the circling particle  $\mu$  (it is called the first adiabatic invariant). A rotating charge is equivalent to a current so we can define its magnetic moment in the following way:

$$\boldsymbol{\mu} = I \, \pi \, \boldsymbol{r}_g \cdot \boldsymbol{n}$$

where the current *I* is given by  $I = \frac{dq}{dt} = \frac{q \Omega_c}{2\pi}$ .

Then we get for  $\mu$ :

$$\boldsymbol{\mu} = \frac{e \,\Omega_c}{2\pi} \pi \frac{v_\perp^2}{\Omega_c^2} = \frac{1}{2} m \frac{v_\perp^2}{B} \boldsymbol{n}$$

This is an adiabatic invariant and is conserved when the length scale of the variation of the magnetic field is significantly larger than the radius of the gyromagnetic motion. Then its invariance can be easily proved. Let's start from the variation of the perpendicular component of the particle velocity:

$$\frac{d}{dt}(mv_{\perp}) = q\left(\boldsymbol{E}_{\perp} + v_{\perp} \times \boldsymbol{B}\right)$$

if we multiply both sides by  $v_{\perp} \perp$  we get:

$$m\frac{\boldsymbol{v}_{\perp} \cdot d\,\boldsymbol{v}_{\perp}}{dt} = \frac{d}{dt}(\frac{1}{2}m\,\boldsymbol{v}_{\perp}^2) = q\,\boldsymbol{v}_{\perp} \cdot \boldsymbol{E}_{\perp}$$

At this point we integrate the previous formula over one complete gyromagnetic period  $T = \frac{2\pi}{\Omega_c}$  to get:

$$\int_0^T \frac{d}{dt} \left(\frac{1}{2} m v_\perp^2\right) dt = \Delta \left(\frac{1}{2} m v_\perp^2\right) = q \int_0^T v_\perp \cdot \boldsymbol{E}_\perp dt = q \oint \boldsymbol{E}_\perp \cdot d\boldsymbol{I}$$

We can apply Faradys' law and express the electric field as the variation of the magnetic field:

$$q \oint \boldsymbol{E}_{\perp} \cdot d\boldsymbol{l} = -\int_{\Sigma} \frac{\partial \boldsymbol{B}}{\partial t} ds$$

Here  $\Sigma$  is the area enclosed by a gyromagnetic orbit. Since we have assumed that the variation of **B** is small over a gyromagnetic radius, than we can approximate the integral on the right as:  $Z \downarrow$ 

$$-\int_{\Sigma} \frac{\partial \boldsymbol{B}}{\partial t} ds = -q \frac{dB}{dt} \cdot \Sigma = q \frac{dB}{dt} \pi r_c^2$$

Why the change in sign from the second to the third expression?



When the line-integral is transformed into a surface integral by the Faradys' law, a convention on the direction of the line integral (anti-clockwise) is adopted (see side figure). However, the line integral is computed along the gyromagnetic motion with is clockwise (for positive charges, than the q changes sign). As a consequence, we have to assume that the normal **n** points along *-z* or, in alternative, that the area is negative. At this point we can substitute the previously computed expression for  $r_c$  to get:

$$\Delta(\frac{1}{2}mv\perp^2) = \Delta\epsilon_{\perp} = \pi q \frac{dB}{dt} \frac{m^2 v_{\perp}^2}{q^2 B^2} = 2\pi \frac{dB}{dt} \frac{\frac{1}{2}mv_{\perp}^2}{B} \frac{m}{qB} = (\frac{2\pi}{\Omega_c})\frac{\epsilon_{\perp}}{B}\frac{dB}{dt} = T\frac{dB}{dt}\frac{\epsilon_{\perp}}{B}$$

If the magnetic field changes smoothly on time because of the motion of the particle along the fiels line (due to  $v_{\parallel}$ ), we can substitute  $T\frac{dB}{dt}$  with  $\Delta B$  so that the Farady's law lead to the following equation:

$$\Delta \epsilon_{\perp} = \frac{\Delta B}{B} \epsilon_{\perp}$$

From calculus we know that if  $\frac{\Delta y}{\Delta x} = \frac{y}{x}$  then :

$$\Delta(\frac{y}{x}) = \frac{\partial(\frac{y}{x})}{\partial x} \Delta x + \frac{\partial(\frac{y}{x})}{\partial y} \Delta y = -\frac{y}{x^2} \Delta x + \frac{\Delta y}{x} = -\frac{y}{x^2} \Delta x + \frac{y}{x^2} \Delta x = 0$$

For this reason  $\Delta \mu = \Delta(\frac{\epsilon_{\perp}}{B}) = 0$  and the magnetic moment is an adiabatic invariant for slow changes in time or small changes in space of the magnetic field. The system has then two constants,  $\mu$  and the kinetic energy  $\epsilon = \epsilon_{\perp} + \epsilon_{\parallel} = \frac{1}{2}m(v_{\perp}^2 + v_{\parallel}^2)$ . When the particle moves towards a region of space where the field lines converge, the value of *B* grows. To keep  $\mu$  constant,  $\epsilon_{\perp} = \frac{1}{2}mv_{\perp}^2$  must increase at the expenses of  $\epsilon_{\parallel} = \frac{1}{2}mv_{\parallel}^2$  until  $v_{\parallel} = 0$ . When this limit is reached, the motion reverses and the particles moves towards regions where the filed lines are widening with increasing parallel velocity.

This is the principle of the magnetic bottle confinement. It works also for charged particles in the dipolar magnetic field of the Earth. For larger values of the magnetic fields the radius  $r_c = \frac{2\mu}{qv_{\perp}}$  decreases while the perpendicular velocity  $v_{\perp} = \sqrt{\frac{2\mu B}{m}}$ increases.



### Van Allen Belts



The Earth's Van Allen Belts consists of highly energetic ionized particles trapped in the Earth's geomagnetic fields. There is a small but very dense inner belt extending from about 1 to 3 Earth radii ( $R_E$ ) with a mximum around 2  $R_E$  consisting mainly of high energy protons (10-50 Mev) mostly generated by the beta decay of neutrons related to cosmic rays (CRAN) and by anomalous cosmic rays (ACR, ions picked up from the interstellar neutral gas at the heliosphere). Heavier ions like  $O^+$  (1-00 Kev) are also present possibly produced by the interaction of the gyromagnetic particles with the upper atmosphere or by solar storms. Low energy  $e^-$  are also

populate the inner belt.

PROTON (E>10 MeV) AND ELECTRON (E>1 MeV) INTEGRAL FLUXES



to solar wind storms and distortions can occur like that shown in the side figure (Baker et al., Nature 432, 878–881, 2004) detected by SAMPEX mission:

The outer belt is made mostly of high-energy  $e^-$  and low energetic protons (that form plasma with  $e^-$ ). It extends from about 3 to 9 Earth radii with the center at about 6 Earth radii. There can be large variations due



# APPENDIX A: PROOF OF THE FORMULA FOR DRIFT MOTION DUE TO FIELD CURVATURE.

Let's start from the case where the gyromagnetic motion of a charged particle is perturbed by an electric field. The configuration of the fields is such that the magnetic field  $\mathbf{B}$  is directed along the z-axis while the  $\mathbf{E}$  field is constant and uniform with any direction. The Lorentz's force has the following expression:

$$m\dot{v} = q(E + v \times B)$$

We introduce a reference frame moving with velocity  $v_{RF} = \frac{E \times B}{B^2}$ . In this reference frame, the velocity of the charged particle is  $u = v - v_{RF} = v - \frac{E \times B}{B^2}$ . Since both the fields **E** and **B** are constants,  $\dot{u} = \dot{v}$ . In this new reference frame, we can study the trajectory of the charged particle since we know its equation of motion:

$$m \dot{\boldsymbol{u}} = m \dot{\boldsymbol{v}} = q \left( \boldsymbol{E} + \boldsymbol{v} \times \boldsymbol{B} \right) = q \left( \boldsymbol{E} + \left( \boldsymbol{u} + \frac{\boldsymbol{E} \times \boldsymbol{B}}{\boldsymbol{B}^2} \right) \times \boldsymbol{B} \right) = q \left( \boldsymbol{E} + \boldsymbol{u} \times \boldsymbol{B} + \frac{(\boldsymbol{E} \times \boldsymbol{B}) \times \boldsymbol{B}}{\boldsymbol{B}^2} \right)$$

Recalling the rules that relate the cross product of 3 vectors to the dot product of the same vectors:

$$(A \times B) \times C = (A \cdot C) B - (B \cdot C) A$$

we get:

$$m \, \boldsymbol{u} = q \left( \boldsymbol{E} + \boldsymbol{u} \times \boldsymbol{B} + \frac{(\boldsymbol{E} \cdot \boldsymbol{B}) \, \boldsymbol{B}}{B^2} - \boldsymbol{E} \right) = q \left( \boldsymbol{u} \times \boldsymbol{B} + (\boldsymbol{E} \cdot \boldsymbol{\hat{b}}) \, \boldsymbol{\hat{b}} \right)$$

where  $\hat{b} = \frac{B}{B} = z$ . Let's split the velocity of the particle in a component  $u_{\parallel} = u \cdot \hat{b}$  parallel to the zaxis and in a component  $u_{\perp} = |u - u_{\parallel} \cdot \hat{b}|$  perpendicular to the z-axis and then to the magnetic field **B**. The equation of motion can be split in two equations for each of the velocity components. For the parallel component we get:

$$m \, \boldsymbol{u} \cdot \boldsymbol{\hat{b}} = m \, \dot{\boldsymbol{u}}_{\parallel} = q \left( (\boldsymbol{\hat{b}} \cdot \boldsymbol{\hat{b}}) (\boldsymbol{E} \cdot \boldsymbol{\hat{b}}) + (\boldsymbol{u} \times \boldsymbol{B}) \cdot \boldsymbol{\hat{b}} \right) = q \, E_{\parallel}$$

The motion along the z-axis is accelerated by the component of the electric field along the z-axis. We can go back to the original reference frame and compute the equation of the velocity  $v_{\parallel}$  since:

$$u_{\parallel} = \boldsymbol{u} \cdot \boldsymbol{\hat{b}} = \boldsymbol{v} \cdot \boldsymbol{\hat{b}} - \frac{(\boldsymbol{E} \times \boldsymbol{b})}{\boldsymbol{\hat{b}}} = \boldsymbol{v} \cdot \boldsymbol{\hat{b}} = v_{\parallel}$$

As a consequence, the equation for the parallel component of the velocity in the initial reference frame is:

$$v_{\parallel} = \frac{q}{m} E_{\parallel} t + v_{\parallel 0}$$

The most interesting part is the motion in the planet perpendicular to the z-axis:

$$m \dot{\boldsymbol{u}} - m \dot{\boldsymbol{u}}_{\parallel} \hat{\boldsymbol{b}} = q \left( \hat{\boldsymbol{b}} \left( \boldsymbol{E} \cdot \hat{\boldsymbol{b}} \right) + \boldsymbol{u} \times \boldsymbol{B} - \boldsymbol{E}_{\parallel} \hat{\boldsymbol{b}} \right) = q \left( \boldsymbol{u} \times \boldsymbol{B} \right) = q \boldsymbol{u}_{\perp} \times \boldsymbol{B} = m \, \dot{\boldsymbol{u}}_{\perp}$$

This equation implies that the motion in the new reference frame is a gyromagnetic motion around the new z-axis. In the initial reference frame it is a gyromagnetic motion with a guiding center which

moves with a velocity  $v_{gc} = v_{\parallel} \hat{b} + \frac{E \times B}{B^2}$ . The first term is a motion along the direction of the filed while the second term is the drift motion in a direction perpendicular to the field **B**. Along the derivation of the equation for  $v_{gc}$  we never used any relation between the magnetic and electric field, Maxwell's equations were never used. This implies that the derivation holds if instead of the Electric field we use any other force field like gravity. In this case the drift velocity would be:

$$\boldsymbol{v}_{gc} = \boldsymbol{v}_{DG} = \frac{m}{q B^2} \boldsymbol{G} \times \boldsymbol{B}$$

since it is possible to make the substitution  $E = \frac{F}{q} = \frac{m}{q}G$  along the proof of the drift velocity. The same holds for the centrifugal force and this leads to the demonstration of the formula for  $v_{DI}$ . If the field lines of the magnetic field **B** are bent, we can introduce a local reference frame with the z-axis along the tangent to the field line. The gyromotion is supposed to follow the field line moving along it with the velocity  $v_{\parallel}$ . Locally, this is a circular motion whose center is located along the perpendicular direction to the local tangent of the field line and whose radius is the local curvature radius  $R_c$ . In a reference frame moving with  $v_{\parallel}$  along the field line the particle feels a centrifugal force equal to:

$$F_{cf} = m \frac{v_{\parallel}^2}{R_c} \hat{\boldsymbol{r}} = m v_{\parallel}^2 \frac{\boldsymbol{R}_c}{R_c}$$

If we assume that this force is felt by a particle performing a gyromotion, than we can resort to the results for the Electric field by substituting  $E = \frac{F_{cf}}{q}$  so that the formula for the drift is:

$$\boldsymbol{v}_{\boldsymbol{D}\boldsymbol{I}} = \frac{\boldsymbol{F}_{cf} \times \boldsymbol{B}}{\boldsymbol{q} \boldsymbol{B}^2} = m \, v_{\parallel}^2 \frac{\boldsymbol{R}_c \times \boldsymbol{B}}{\boldsymbol{R}_c \boldsymbol{q} \boldsymbol{B}^2} = m \, v_{\parallel}^2 \frac{\boldsymbol{B} \times \boldsymbol{n}}{\boldsymbol{R}_c \boldsymbol{q} \boldsymbol{B}^2}$$

dove  $R_c = -n$ .

# Magnetosphere of the Earth.

The interaction of the solar wind with the magnetic field of the Earth gives origin to a very complex environment called the magnetosphere (see figure3 below).



The radial solar wind is deflected by the magnetic field lines and a complex structure is created around the planet characterized by complicate plasma fluxes. We can derive an approximate estimate of the average size of the magnetosphere and how it responds to changes in the solar wind parameters like density and average velocity. We first estimate the pressure due to the solar wind particles following the lines of the particle-in-a-box approach for the computation of a ideal gas pressure. The variation of the momentum due to a single particle of the solar wind plasma interacting with an imaginary surface  $dx \cdot dy$  perpendicular to the Sun-Earth direction (*z*-axis) is:

$$\Delta p = -mv$$

Usually, in the kinetic theory for ideal gas  $\Delta p = -2 mv$  since it is assumed that the gas particle bounces back on the same direction. In our case the particle is *deflected* and we can assume as a first approximation that the deflection occurs at 90° respect to the original velocity. As a consequence, the variation of momentum in the radial direction after the interaction is only  $\Delta p = -mv$  since the motion after the interaction is in a direction perpendicular to the initial motion. The total force due to N particles interacting with the surface (N is the number of particles reaching the surface in a time dt) is then:

$$F = \frac{\Delta p \cdot N}{dt} = \frac{N m v^2}{dz}$$

The pressure is the force divided by the area of the surface so:

$$P = \frac{F}{A} = \frac{N m v^2}{dx \cdot dy \cdot dz} = \frac{N m v^2}{V} = \rho v^2$$

The magnetic pressure which is the force of deflection (or radius of the induced gyromotion) is expressed by the formula:

$$P_{M} = \frac{B^{2}}{2\mu_{0}}$$

Equating the two expression for the pressure we find the radius were the two forces balances:

$$\rho v^2 = \frac{B^2}{2 \mu_0} = \frac{M_B^2}{2 \mu_0 r^6}$$

where MB is the magnetic dipole moment for the Earth. Solving the above equation in r we get an approximate value for the magnetosphere radius:

$$r_{M} = \left(\frac{M_{B}^{2}}{2\mu_{0}\rho v^{2}}\right)^{\frac{1}{6}}$$

For average values of the solar wind density ( $\rho \approx 5 \text{ p}^+ \text{ per cm}^3$ ,  $v \approx 300 \text{ km/s}$ ) we get  $r_M \approx 10 \text{ R}_E$ .